

Amateur Radio Astronomy

Projet ART ***Awesome Radio Telescope***




Ruben Barbosa
2016

FOTO: © Miguel Claro | ESO / ALMA


How to explore the Universe?

The search for ET life follows 3 main directions:

1. looking for **primitive life** (bacteria) → the **solar system**, by sending automated interplanetary probes,
2. **biomarker** detection (ozone) → in **exoplanets** similar to Earth located in habitable zones and
3. **complex life** demand (iET) → forms of life with technology similar to ours, by SETI




The rise of ART Project



NASA/SETI/JPL

The electromagnetic spectrum



Thermal Imaging



ART Project - OAPBG

If our eyes were sensitive to radio ...


- ✓ The sky was black with cosmic background radiation
- ✓ There would be no stars visible
- ✓ Day and night have the same appearance
- ✓ Our Sun would be a very bright disk
- ✓ The moon would appear as a homogeneous disk, without craters
- ✓ At 21 cm would observe the Milky Way as a band across the sky



false colors

Microwave window

- ✓ Radio broadcast, covers frequencies from a few MHz (100 m) to frequencies of 300 GHz (1 mm)
- ✓ **Microwave window** 1 GHz to 10 GHz (30 cm to 3 cm), characterized by low background noise
- ✓ **Minimum** frequency limit is caused by the strong absorption in the atmosphere of the elements: O_2 , CO_2 , H_2O
- ✓ **Maximum** frequency limit is due to the opacity of the ionosphere



Emission sources


✓ Thermal radiation:

- ✓ Black body: temperature of the planets
- ✓ Bremsstrahlung (free-free): HII in M42
- ✓ Spectral Ringer HI 21cm in the Milky Way

✓ Non-thermal radiation:

- ✓ Synchrotron: magnetic fields in pulsars, supernovas, radio galaxies

Spectral distribution of general energy




Bremsstrahlung (free-free): HII in M42



Carlos Soares / OAPBG


Solar System radio

The universe changes when seen in band radio ...



NRAO/AUI

Radio galaxy Cygnus A




NRAO/AUI

Milky Way in the optical and radio (408 MHz)


Optical

A. Mellinger Photomosaic



Radio Continuum (408 MHz)


Bonn, Jodrell Bank, and Parkes



NASA


Atomic Hydrogen

21 cm Dickey-Lockman




Molecular Hydrogen

115 GHz Columbia-GISS




Infrared

12, 60, 100 μ m IRAS




Near Infrared

1.25, 2.2, 3.5 μ m COBE/DIRBE



Gamma Ray

>100MeV CGRO/EGRET




Radiation mediation methods

✓ On/off-source

The source is measured for a few minutes and then the antenna is pointed at a site sky (cold) during the same period of time, where the source will pass again.

✓ Transit

The telescope is targeted to the area where the source will carry over (2 hours prior to measuring the background radiation).



NRAO/ Ian Parker

Scientific importance of radio observing



- ✓ Cosmic microwave background radiation (Penzias and Wilson, 1963)

- ✓ Spiral structure of the Milky Way and rotation curve (note the letter of 21 cm of hydrogen)



- ✓ Discovery of new objects: Pulsars (Antony Hewish, 1974) and Quasars

- ✓ Observation of molecular clouds and star formation zones



NASA, APOD 2008-03-10, Graeme L. White & Glen Cozens

Resolution comparison

- ✓ Telescope Effelsberg radio (D = 100 m)
- ✓ Telescope in optical (D = 1 m)

$$\alpha = \frac{1,22 \lambda}{D}$$

$$\lambda_{\text{optical}} \sim 600 \text{ nm}$$

$$\lambda_{\text{radio}} \sim 6 \text{ mm}$$

α_{radio}


α_{optical}

- ✓ The resolving power of an optical telescope in (1 m diameter) is about 100 times higher than that achieved with one of the largest steerable radio telescopes.

- ✓ How to increase the resolving power? Interferometry.

MPIfR (Norbert Junkes)


Scheme ART



A strong source has intensity few Jy
A weak source has intensity few mJy


Why 11,2 GHz?

- ✓ Low cost of material (wide availability of material for satellite TV)
- ✓ Acceptable resolving power
- ✓ Minimum level of sky noise
- ✓ Strong immunity to artificial noise (can be used in urban areas)
- ✓ Radio astronomical sources in this frequency are not numerous or powerful, which means that small satellite dishes (up to 1.5 m) allow only observe the Sun (thermal component) and the moon, and
- ✓ Geostationary satellites can be a source of artificial noise.



Andrew Clegg/National Science Foundation

ART's cockpit (Mars detection in opposition)



Global definitions

Frequency (f)	11.200 MHz
Wavelength (λ)	2,677 cm
Diameter of the aperture (D)	3,00 m
Antenna aperture efficiency (η)	0,50
Antenna depth (d) / Focal lenght (z)	30 m (z = 1,88 m)

Analysys parabolic reflector antenna with circular symmtery

Antenna gain (Ga_max)	61.988 times
Antenna gain (Ga_max_dB)	47,9 dB
Effective diameter of the antenna (Deff)	2,12 m (Aeff=3,534 m2)
Theoretical Beamwidth - HPBW (θ) = $\sim 0,7439^\circ$	0,744 $^\circ$ = 44,64 arcmin
Theoretical Beamwidth (max) - HPBW (λ/D)	0,624 $^\circ$ = 37,42 arcmin
Diameter of the full covered area (BWFN)	1,49 $^\circ$ (+3dB HPBW)
Geometric area of the dish (A)	7,07 m2
Primus focus (0,32 - 0,43)	1,1 m

Receiver parameters

GLNB	55 dB
Line amplifier gain + GIF	0 dB
Losses coaxial cable and connectors	1 dB
Total gain (G)	54,0 dB
Total gain (G)	251.189 times
LNB noise (Fr_dB)	0,4 dB
Noise temperatura of the receiver (Tr)	28,9 k
Integration time (τ)	0,1 ' (RMS10%=3,8 ')
Bandwidth of the receiver (BW)	250 MHz
Impedance F connector for RF-IF input (R)	75 ohm
Voltage input to the doide detector	1,37 mV ($\Delta dBm =$,)

Parameters of the observed radio source

Fictitious angular diameter - non thermal	2,81 arcmin
Apparent angular diameter	0,30 arcmin
Solid angle subtented by the source (Ω_r)	5,98E-09 rad^2
RT's acceptance (Ω)	5,30E-04 rad^2
Real flow of the radio source (Sv)	5,1 Jy
Brightness temperature (ΔT)	221 k
Empirically adjustment (A)	3,59654

Flux correction

Measured temperature	210 k
Source measured flux	4,8 Jy
Atmospheric opacity (τ_a)	0,05
Elevation angle of the antenna (ϕ)	70,0
Zenith opacity estimate (τ_z)	0,05
Elevation to predict	85,0 $^\circ$
Flux density estimate	4,9 Jy (T = 211 k)

Antenna temperature


BT of the cosmic μ wave background (Tcmb)	2,8 k
BT of the atmosphere with attenuation (Tatm)	
BT of the atmosphere (Tatm)	3,6 k
Noise of the ground (Tgnd)	4,6 k ($\phi=1,498^\circ$)
Azimuthal distribution of BT of the RS $\rightarrow T_s(\phi)$	221,4 k ($\phi=0,000^\circ$)
Sky temp.: T(0) T(out) $\Delta=221,4$ k	232,4 k (T = 11,0 k)
Antenna temp.: Ta(0) Ta(out) $\Delta=0,009$ k	0,087 k (T = 0,077 k)
System temp.: Tsys(0) Tsys(out) $\Delta=0,009$ k	29,0 k (T = 29,0 k)

Sensitivity of the receiver

Minimum measurable change (Δt_{min})	0,0058 k
Min. detectable source flux density (ΔS_{min})	4,533 Jy
Coefficient of performance degradation (ξ)	5
Detection signal variation	0,000 k




Drifting versus HPBW

Source drift time (BWFN)	1110 "
Source transit angle (BWFN)	4,63 $^\circ$
Source drift angle from equator	#NÚM! $^\circ$
Source declination (δ)	22,0 $^\circ$
Max. time drift (HPBW)	161,4 " = 3 '
HPBW	4,288 $^\circ$





Theoretical models

Antenna gain pattern




Green Bank Image Gallery

Scope of observations



Software: Radio-Eyes



Can we communicate with iET?

- ✓ The next region of the spectrum of 1420 MHz (21 cm), radio broadcasts have very low noise levels and absorption by the interstellar medium is minimal.
- ✓ Hydrogen being one of the fundamental building blocks of the universe, one o'clock ET civilization will also possess this knowledge.
- ✓ Civilizations ET distance of 150 al equipped with RT 300 meters similar to Arecibo, can transmit radio signals detectable on Earth.
- ✓ Terrestrial antennas 10 meters can reach ~ 21,500 stars (600 s).
- ✓ Terrestrial antennas 3 meters have an average range of 5 star ...

Global definitions	
Frequency (f)	1.420 MHz
Wavelength (λ)	21,11 cm
Diameter of the aperture (D)	10,00 m
Antenna aperture efficiency (η)	0,75
Integration time (τ)	600,0
Bandwidth of the receiver (BW)	256 MHz
SETI	
ETI transmitter power (P)	10.000.000 Mw
Diameter of the aperture antenna ETI	300 m
Distance from Earth (R)	46,8 pc
Stars in our range	21.479

Radio Telescope & Milky Way
by Seth McCubbin

Intelligent extraterrestrial life

Establishing contact ...

A few imitation gestures

DNA Hybridization ~ 1%

Also my 3 year old son ...
I do mental calculations of astrophysics

DNA Hybridization ~ 1%

Also my 3 year old son ...

Communicational interest?

Questions we would like to make a iET?



Communicational interest?

1. Is there any creator of the universe?
Big Bang?
2. What is the universe made? Dark matter? Dark Energy?
3. What is the purpose of the universe?
Real? Simulation?
4. Are we alone in the Galaxy?

The End



Rahi Varsani